ELSEVIER

Biomembranes

Vol. 1238, 1995

BBA Biochimica et Biophysica Acta

Author Index

Akutsu, H., see Shin, K.-h. (1238) 42
Antunes-Madeira, M.C., see Videira, R.A. (1238) 22

Bally, M.B., see Webb, M.S. (1238) 147
Borges, C.P.F., see Borissevitch, I.E. (1238) 57
Borissevitch, I.E., Borges, C.P.F., Yushmanov, V.E. and Tabak, M.
Localization of dipyridamole molecules in ionic micelles: effect of micelle and drug charges (1238) 57

Castaing, M., see Loiseau, A. (1238) 107
Chang, Y.-F. and Charles, A.K.
Uptake and metabolism of Δ^1-piperidine-2-carboxylic acid by synaptosomes from rat cerebral cortex (1238) 29
Charles, A.K., see Chang, Y.-F. (1238) 29
Ciampolillo-Bates, F., see Karlson, K.H. (1238) 197
Custódio, J.B.A., see Videira, R.A. (1238) 22

Dalla Serra, M., see Menestrina, G. (1238) 72
De Kruijff, B., see Speelmans, G. (1238) 137
De Wolf, F.A., see Speelmans, G. (1238) 137
Dietrich, C. and Tampé, R.
Charge determination of membrane molecules in polymer-supported lipid layers (1238) 183

Fujiwara, T., see Shin, K.-h. (1238) 42
Furblur, U.G., see Lear, J.D. (1238) 34

Gogarten, J.P., see Hilario, E. (1238) 94
Goncz, K.K. and Rothman, S.S.
A trans-membrane pore can account for protein movement across zymogen granule membranes (1238) 91
Grøndahl, M.L., see Munck, L.K. (1238) 49

Halestrap, A.P., see Jackson, V.N. (1238) 193
Hartsel, S.C., see Wolf, B.D. (1238) 156
Herick, K. and Krämer, R.
Kinetic and energetic characterization of solute flux through the reconstituted aspartate/glutamate carrier from beef heart mitochondria after modification with mercurials (1238) 63
Hilario, E. and Gogarten, J.P.
The V-ATPase *A* subunit gene (*vma-1*) from *Giardia lamblia* (1238) 94
Hill, M., see Loiseau, A. (1238) 107
Hugo, F., see Menestrina, G. (1238) 72

Irimura, T., see Koike, C. (1238) 99

Jackson, V.N., Price, N.T. and Halestrap, A.P.
cDNA cloning of MCT1, a monocarboxylate transporter from rat skeletal muscle (1238) 193
Jing, N. and Urry, D.W.
Ion pair binding of Ca^{2+} and Cl^- ions in micellar-packaged gramicidin A (1238) 12
Jing, N., Prasad, K.U. and Urry, D.W.
The determination of binding constants of micellar-packaged gramicidin A by ^{13}C-and ^{23}Na-NMR (1238) 1

Kakiuchi, T., see Koike, C. (1238) 99
Karlson, K.H., Ciampolillo-Bates, F., McCoy, D.E., Kizer, N.L. and Stanton, B.A.
Corrigendum to "Cloning of a cGMP-gated cation channel from mouse kidney inner medullary collecting duct" [Biochim. Biophys. Acta 1236 (1995) 197–200] (1238) 197
Kizer, N.L., see Karlson, K.H. (1238) 197
Koike, C., Oku, N., Watanabe, M., Tsukada, H., Kakiuchi, T., Irimura, T. and Okada, S.
Real-time PET analysis of metastatic tumor cell trafficking in vivo and its relation to adhesion properties (1238) 99
Krämer, R., see Herick, K. (1238) 63
Kubalski, A.
Generation of giant protoplasts of *Escherichia coli* and an inner-membrane anion selective conductance (1238) 177

Lally, E.T., see Lear, J.D. (1238) 34
Le, H.-b., see Urbina, J.A. (1238) 163
Lear, J.D., Furblur, U.G., Lally, E.T. and Tanaka, J.C.
Actinobacillus actinomycetemcomitans leukotoxin forms large conductance, voltage-gated ion channels when incorporated into planar lipid bilayers (1238) 34
Loiseau, A., Hill, M., René-Corail, L. and Castaing, M.
Transport of competing Na and K ions by (222)C_{10}-cryptand, an ionizable mobile carrier: effects of pH and temperature (1238) 107

Madeira, V.M.C., see Videira, R.A. (1238) 22
Maeda, H., see Shin, K.-h. (1238) 42
May, J.M., Qu, Z.-c. and Whitesell, R.R.
Ascorbate is the major electron donor for a transmembrane oxidoreductase of human erythrocytes (1238) 127
Mayer, L.D., see Webb, M.S. (1238) 147
McCoy, D.E., see Karlson, K.H. (1238) 197
Menestrina, G., Ropele, M., Dalla Serra, M., Pederzolli, C., Hugo, F., Pellet, S. and Welch, R.A.
Binding of antibodies to functional epitopes on the pore formed by *Escherichia coli* hemolysin in cells and model membranes (1238) 72
Montez, B., see Urbina, J.A. (1238) 163
Munck, L.K., Grøndahl, M.L. and Skadhauge, E.
Beta-amino acid transport in pig small intestine in vitro by a high-affinity, chloride-dependent carrier (1238) 49

Neamtu, S., see Turcu, I. (1238) 81

Okada, S., see Koike, C. (1238) 99
Okada, S., see Oku, N. (1238) 86
Oku, N., Tokudome, Y., Tsukada, H. and Okada, S.
Real-time analysis of liposomal trafficking in tumor-bearing mice by use of positron emission tomography (1238) 86
Oku, N., see Koike, C. (1238) 99
Oldfield, E., see Urbina, J.A. (1238) 163

Patterson, J., see Urbina, J.A. (1238) 163
Pederzolli, C., see Menestrina, G. (1238) 72
Pekerar, S., see Urbina, J.A. (1238) 163
Pellet, S., see Menestrina, G. (1238) 72

Prasad, K.U., see Jing, N. (1238) 1
Price, N.T., see Jackson, V.N. (1238) 193

Qu, Z.-c., see May, J.M. (1238) 127

René-Corail, L., see Loiseau, A. (1238) 107
Ropele, M., see Menestrina, G. (1238) 72
Rothman, S.S., see Goncz, K.K. (1238) 91

Shin, K.-h., Maeda, H., Fujiwara, T. and Akutsu, H.
Molecular miscibility of phosphatidylcholine and phosphatidylethanolamine in binary mixed bilayers with acidic phospholipids studied by ^{2}H- and ^{31}P-NMR (1238) 42
Skadhauge, E., see Munck, L.K. (1238) 49
Slotte, J.P.
Lateral domain heterogeneity in cholesterol/phosphatidylcholine monolayers as a function of cholesterol concentration and phosphatidylcholine acyl chain length (1238) 118
Speelmans, G., Staffhorst, R.W.H.M., De Wolf, F.A. and De Kruijff, B.
Verapamil competes with doxorubicin for binding to anionic phospholipids resulting in increased internal concentrations and rates of passive transport of doxorubicin (1238) 137
Staffhorst, R.W.H.M., see Speelmans, G. (1238) 137
Stanton, B.A., see Karlson, K.H. (1238) 197

Tabak, M., see Borissevitch, I.E. (1238) 57
Tampé, R., see Dietrich, C. (1238) 183
Tanaka, J.C., see Lear, J.D. (1238) 34
Tokudome, Y., see Oku, N. (1238) 86
Tsukada, H., see Koike, C. (1238) 99
Tsukada, H., see Oku, N. (1238) 86
Turcu, I. and Neamtu, S.
Dimensional distribution of human erythrocytes obtained from electropermeabilization experiments (1238) 81

Urbina, J.A., Pekerar, S., Le, H.-b., Patterson, J., Montez, B. and Oldfield, E.
Molecular order and dynamics of phosphatidylcholine bilayer membranes in the presence of cholesterol, ergosterol and lanosterol: a comparative study using ^{2}H-, ^{13}C- and ^{31}P-NMR spectroscopy (1238) 163
Urry, D.W., see Jing, N. (1238) 1
Urry, D.W., see Jing, N. (1238) 12

Videira, R.A., Antunes-Madeira, M.C., Custódio, J.B.A. and Madeira, V.M.C.
Partition of DDE in synthetic and native membranes determined by ultraviolet derivative spectroscopy (1238) 22

Watanabe, M., see Koike, C. (1238) 99
Webb, M.S., Wheeler, J.J., Bally, M.B. and Mayer, L.D.
The cationic lipid stearylamine reduces the permeability of the cationic drugs verapamil and prochlorperazine to lipid bilayers: implications for drug delivery (1238) 147
Welch, R.A., see Menestrina, G. (1238) 72
Wheeler, J.J., see Webb, M.S. (1238) 147
Whitesell, R.R., see May, J.M. (1238) 127
Wolf, B.D. and Hartsel, S.C.
Osmotic stress sensitizes sterol-free phospholipid bilayers to the action of Amphotericin B (1238) 156

Yushmanov, V.E., see Borissevitch, I.E. (1238) 57